


Atomic Force Microscope (AFM)

The Atomic Force Microscope



More than 50 AFM images are on this website. We have used AFM to characterize extensively nucleosomes and chromatin fibers.

Selected references:

Leuba, S. H., Yang, G., Robert, C., Samori, B., van Holde, K., Zlatanova, J. and Bustamante, C. Three-dimensional structure of extended chromatin fibers as revealed by tapping-mode scanning force microscopy. Proc. Natl. Acad. Sci. USA **91**, 11621-11625, 1994.

Leuba, S. H., Bustamante, C., Zlatanova, J. and van Holde, K. Contributions of linker histones and histone H3 to chromatin structure: Scanning force microscopy studies on trypsinized fibers. Biophys. J. **74**, 2823-2829, 1998.

Leuba, S. H., Bustamante, C., van Holde, K. and Zlatanova, J. Linker histone tails and N-tails of histone H3 are redundant: scanning force microscopy studies of reconstituted fibers. Biophys. J. **74**, 2830-2839, 1998.

Leuba, S. H. and Bustamante, C. Analysis of Chromatin by Scanning Force Microscopy. In Methods in Molecular Biology, Becker, P., Ed. **119**, 143-160, 1999.

Karymov, M. A., Tomschik, M., Leuba, S. H., Caiafa, P. and Zlatanova, J. DNA Methylation-dependent chromatin fiber compaction in vivo and in vitro: Requirement for linker histone. *FASEB J.* **15**, 2631-2641, 2001.

Tomschik, M., Karymov, M. A., Zlatanova, J. and Leuba, S. H. The archaeal histone-fold protein HMf organizes DNA into bona fide chromatin fibers. *Struct. Fold. Des.* **9**, 1201-1211, 2001.

An, W., Palhan, V. B., Karymov, M. A., Leuba, S. H., and Roeder, R. G. Selective requirements for histone H3 and H4 N-Termini in p300-dependent transcriptional activation from chromatin. *Mol. Cell* **9**, 811-821, 2002.